



## Immunization Schedule with Combination Vaccines



Schedule
Using
PENTACEL®5
&
KINRIX™











HepA

KINRIX<sup>™5</sup>

DTaP, IPV

MMR

**Varicella** 

**EVERY FALL: FLU VACCINE<sup>4</sup> Children age 6 months through 18 years** 

**EVERY FALL: FLU VACCINE<sup>4</sup> Children age 6 months through 18 years** 

## Combination vaccines are not identical.

Make sure the vaccine you administer contains the antigens on the doctor's order. Keep it simple, stick with the same product.

This is a suggested schedule for VFC providers utilizing combination vaccines. For alternatives and details, consult the latest "Recommended Immunization Schedules for persons aged 0-6 years and 7-18 years, United States." For more info visit www.eziz.org

- <sup>1</sup> A dose of Hepatitis B vaccine is not necessary at 4 months if doses are given at birth and 2 months but may be included as part of a combination vaccine.
- 2 The rotavirus vaccines differ in number of doses and timing. Check the package insert.
- <sup>3</sup> This six month Hib dose is not indicated if Pedvax HIB® and/or COMVAX® are used exclusively for the 2 and 4 month infant doses.
- <sup>4</sup> Influenza vaccine is available in thimerosal-free options. See California Health and Safety Code § 124172.
- <sup>5</sup> Licensed by FDA for children 4 through 6 years with previous doses of INFANRIX™ or PEDIARIX™. ACIP recommends that, whenever feasible, the same manufacturer's DTaP vaccines be used for each dose in the series; however, vaccination should not be deferred because the type of DTaP previously administered is unavailable or unknown. See www.cdc.gov/mmwr/preview/mmwrhtml/mm5739a4.htm

